

On the Fixing of the κ Gauge Symmetry on AdS and Flat
Background:
the Lightcone Action for the Type IIb String on $AdS_5 \otimes S_5$.

Igor Pesando ¹

Dipartimento di Fisica Teorica , Università di Torino, via P. Giuria 1, I-10125
Torino, Istituto Nazionale di Fisica Nucleare (INFN) - sezione di Torino, Italy

ABSTRACT

We explore all the possible ways of fixing the κ symmetry for both branes and strings by means of a constant projector. We find that they can be classified according to their behaviour under Dirac conjugation and conjugation. This latter controls the maximum power of the fermionic variables in which the (super)vielbein can be expanded while the former determines which states cannot be described in the gauge. In particular there exists an interesting class of projectors for which vielbein are at most quadratic in the fermionic variables. As an example we compute the action for the type IIb on a $AdS_5 \otimes S_5$ background with a lightcone-like projector; this action reduces to the usual lightcone GS string action in the flat limit.

¹e-mail: ipesando@to.infn.it, pesando@alf.nbi.dk

1 Introduction.

During the last year there has been a lot of work in finding the action for both $M2$, $D3$ branes ([1],[7],[6]) and the type IIB superstring ([9],[3],[1],[4],[5]) propagating on a $AdS_{p+2} \otimes S_{D-p-2}$ backgrounds. All these actions have been derived using either the supercoset approach ([1]) with the κ symmetry fixed by the so called killing gauge ([2]) or the supersolvable algebra approach ([7]) which automatically yields a natural way to fix the κ symmetry. It turns out that in the cases where the actions have been worked out the two methods actually give the same result since the killing gauge and the supersolvable gauge fix the κ symmetry in the same way.

It is purpose of this letter to explore all the possible way of fixing the κ symmetry by a constant projector of the type

$$P = \frac{1 + \Gamma}{2} \quad \Gamma^2 = 1$$

and discuss the advantages/disadvantages of every possible class of gauge fixings. In order to do this we classify all the possible Γ 's according to its properties under charge conjugation and under Dirac conjugation. It turns out that the latter property is connected with the equation which describes the string states which cannot be described while the former controls the maximum power of the fermionic variables in the κ gauge fixed expression for the supervielbein (either 2 or 16).

In the case of type IIB superstring propagating on a $AdS_5 \otimes S_5$ we determine explicitly the equation for physical states which can be described by the projectors which have two as the maximum power in the fermionic variables expansion for the (super)potentials and we argue that a similar result is valid for all the other cases.. Because of the similarity with the flat case, to which it reduces in the infinite radius limit, we construct the action in the gauge given by the choice $\Gamma = \gamma_0 \gamma_3$.

2 Classifications of the constant projectors.

As the first step we want to classify all the possible constant projectors according to two properties which will turn out to be very important in determining the form of the κ gauge fixed fields and which physical states which can be described.

Here an in the following we assume that Ξ are the fermionic coordinates of the $AdS_{p+2} \otimes S_{D-p-2}$ superspace which transform as spinor *only* under AdS_{p+2} tangent frame rotations and as a scalar for S_{D-p-2} rotations: this is typical of the Kaluza-Klein way of decomposing the superfields and it is different from the supercoset framework where the fermionic coordinates transform as spinor for both AdS_{p+2} and S_{D-p-2} . Since the two approaches have to be equivalent all the results in this letter can be in principle reformulated in the supercoset approach.

We take the projector P on the Ξ coordinates which are to be set to zero in order to fix the κ symmetry to be written as

$$P = \frac{1 + \Gamma}{2} \quad \Gamma^2 = 1 \quad (2.1)$$

and its complement to be $Q = \frac{1 - \Gamma}{2}$. One could wonder why this fixes the κ symmetry. A naive answer is that it throws away half the fermionic dof as it should. A better answer can be only given a posteriori when we can verify that the fermionic kinetic term is not singular or when we can find an explicit κ transformation which moves a generic configuration into one which satisfies this gauge.

For a generic projector P we define the following shorthand notations:

$$\begin{aligned} \bar{P} &= \gamma_0 P^\dagger \gamma_0 \\ P_c &= C(\bar{P})^T C^{-1} \end{aligned}$$

where C , the charge conjugation matrix, and the γ s are the proper matrices for the given spacetime, i.e. they are the AdS γ and C matrices. We define similar shorthand for the Q projector.

We can now introduce the classification according to the behaviour under the Dirac conjugation:

$$Type L : \quad \bar{P} = P \quad \bar{Q} = Q \quad (2.2)$$

$$Type S : \quad \bar{P} = Q \quad \bar{Q} = P \quad (2.3)$$

As we show later in section (3) this classification has to do with whether or not the left invariant forms obtained from a specific coset elements representation are quadratic in the fermionic variables (S) or up to the sixteenth power (L).

We also divide the projectors according to their behaviour under charge conjugation:

$$Type I : \quad P_c = P \quad Q_c = Q \quad (2.4)$$

$$Type II : \quad P_c = Q \quad Q_c = P \quad (2.5)$$

This subdivision, as we will see in section (4), has to do with which physical states can be described by the gauge enforced by the projector P . This will be explicitly proved in the case of the string propagating on $AdS_5 \otimes S_5$ but we believe it is true for all the cases.

Using the conventions of app. A we can now make a series of tables with all the possible Γ defining the projectors up to Lorentz rotations in the world volume of the Dp brane or in its transverse space when we think to $AdS_{p+2} \otimes S_{D-p-2}$ as the near horizon geometry:

² In 10D the matrix Γ has to satisfy another requirement: $[\Gamma, \Gamma_{11}] = 0$ since the fermionic space time coordinates are chiral.

- $AdS_3 \otimes S_3$ case:

	I	II
S	$i\gamma_2, i\gamma_1$	
L		γ_0

where the case $\Gamma = i\gamma_2$ corresponds to the supersolvable algebra;

- $AdS_5 \otimes S_5$ case:

	I	II
S	$\gamma_0\gamma_3, \gamma_0\gamma_4$	$i\gamma_3, i\gamma_4$
L	γ_0	$i\gamma_2\gamma_3, i\gamma_3\gamma_4$

We notice that the choice $\Gamma = i\gamma_4$ corresponds to the supersolvable gauge;

- Flat 10D case:

	I	II
S	$\Gamma_0\Gamma_9$	$i\Gamma_0\Gamma_1\Gamma_2\Gamma_3$
L	$\Gamma_6\Gamma_7\Gamma_8\Gamma_9$	$i\Gamma_8\Gamma_9$

Similar considerations can be easily done for branes and membranes.

3 Short and Long left invariant forms expansion.

We now want to discuss explicitly the meaning of the S and L projectors. To this purpose we write down an explicit representation for the coset elements from which a nice expressions for the fermionic dependency of left invariant forms can be derived. In particular we take as a representative for coset elements the following expression ([8])

$$L(x, \Xi) = L_F(\Xi)L_B(x) \quad (3.1)$$

where

$$L_F = \exp \left(\begin{array}{cc} & (P\Xi)_{\alpha N} \\ (\overline{P\Xi})^{\beta M} & \end{array} \right) \exp \left(\begin{array}{cc} & (Q\Xi)_{\alpha N} \\ (\overline{Q\Xi})^{\beta M} & \end{array} \right) \quad (3.2)$$

where α, β are AdS spinorial indices which label the rows of the upper right block and N, M are column indices on which the R symmetry acts.

We notice that even before fixing the κ symmetry there is a big difference between the projectors S and L , in fact type S projectors yield an expression for L_F quartic in Ξ when explicitly evaluated:

$$G_F = \left(\begin{array}{cc} (1 + \frac{1}{2}P\Xi^R\overline{\Xi}_RQ)_{\alpha}^{\cdot\beta} & (P\Xi)_{\alpha N} \\ (\overline{\Xi}Q)^{\beta M} & 1 \end{array} \right) \left(\begin{array}{cc} (1 + \frac{1}{2}Q\Xi^R\overline{\Xi}_RP)_{\alpha}^{\cdot\beta} & (Q\Xi)_{\alpha N} \\ (\overline{\Xi}P)^{\beta M} & 1 \end{array} \right) \quad (3.3)$$

The same kind of computation gives a very different result for the type L projectors since the equivalent of eq. (3.2) would read

$$G_F = \exp \left(\begin{array}{cc} & (P\underline{\Xi})_{\alpha N} \\ (\underline{\Xi}P)^{\beta M} & \end{array} \right) \exp \left(\begin{array}{cc} & (Q\underline{\Xi})_{\alpha N} \\ (\underline{\Xi}Q)^{\beta M} & \end{array} \right) \quad (3.4)$$

which is a product of two $O(\underline{\Xi}^{16})$ (in the case of maximal supersymmetry) matrices. Because of this complexity we will not consider anymore the projectors of type L.

These properties imply directly the maximum power of the fermionic coordinates in the supervielbeins expansion through the formula $L^{-1}dL = \text{supervielbein} + \text{spin} - \text{connection}$; in particular it is immediate to verify using the κ gauge fixed version of eq. (3.3), i.e. setting $P\underline{\Xi} = 0$, that the κ gauge fixed supervielbeins obtained using S projectors are *at most quadratic* in the fermionic coordinates.

4 The equation for the states which can be described by the κ gauge fixed action.

In this section we concentrate on the consequences of choosing either a type I or a type II projector in the specific case of type IIB superstring on $AdS_5 \otimes S_5$. The same conclusions can essentially be reached in the flat superstring case because the gravitino transformation rule under κ symmetry is essentially eq. (4.3) without the second rhs term which corresponds to the internal S_5 contribution. Moreover we believe that similar conclusions can be drawn for the other cases because while the gravitino transformation rules have additional terms in the κ symmetry expression (4.3) the essential term is given by the first term in rhs which is common to all the cases.

We start writing the first order action for the GS superstring propagating on $AdS_5 \otimes S_5$ in order to fix the notations which we keep almost as in ([9]):

$$\begin{aligned} S = & \frac{T}{2} \int_{\Sigma} \Pi_{\underline{\alpha}}^{\hat{a}} i_* V^{\hat{b}} \wedge e^{\underline{\beta}} \eta_{\hat{a}\hat{b}} \epsilon_{\underline{\alpha}\underline{\beta}} - \frac{1}{4} \Pi_{\underline{\alpha}}^{\hat{a}} \Pi_{\underline{\beta}}^{\hat{b}} \eta_{\hat{a}\hat{b}} \eta^{\underline{\alpha}\underline{\beta}} \epsilon_{\underline{\gamma}\underline{\delta}} e^{\underline{\gamma}} \wedge e^{\underline{\delta}} \\ & + \frac{1}{4} e^{i\phi} i_* (V_+ * A) - \frac{1}{4} e^{-i\phi} i_* (V_- * A) \end{aligned} \quad (4.1)$$

where $e^{\underline{\alpha}}$ ($\underline{\alpha}, \underline{\beta}, \dots \in \{0, 1\}$ worldsheet flat indices) are the worldsheet zweibein, $\Pi_{\underline{\alpha}}^{\hat{a}}$ ($\hat{a}, \hat{b}, \dots \in \{0, \dots, 9\}$ flat spacetime indices) are auxiliary 0-forms related by the equations of motion to $V^{\hat{a}}$, the ten dimensional zehnbein, by $i_* V^{\hat{a}} = \Pi_{\underline{\alpha}}^{\hat{a}} e^{\underline{\alpha}}$ where i_* is the pullback on the string worldsheet Σ due to the superimmersion $i : \Sigma \rightarrow AdS_5 \otimes S_5$. Moreover we have $V_{\pm} * A = \epsilon_{\alpha\beta} V_{\pm}^{\alpha} A^{\beta}$ ($\alpha, \beta \in \{0, 1\}$ $SU(1, 1)$ fundamental irrep indices) with V_{\pm}^{α} the $SU(1, 1)$ scalars and A^{α} the complex 2-forms; finally ϕ is an arbitrary angle we can reabsorb redefining the complex fields A^{α} .

The zweibein equation of motion gives the Virasoro constraints:

$$\frac{\delta S}{\delta e} = 0 \implies \Pi_{\underline{\alpha}} \bullet \Pi_{\underline{\beta}} = \frac{1}{2} \eta_{\underline{\alpha}\underline{\beta}} \Pi^2 \quad (4.2)$$

This action has a κ symmetry which is generated by the action of the Lie derivative with respect to a vector field $\vec{\epsilon}$ which is defined by the properties

$$i_{\vec{\epsilon}} \psi_N = \epsilon_N = \Pi_{\underline{\alpha}}^a \gamma_a \kappa_{\underline{N}}^{\underline{\alpha}} - i \Pi_{\underline{\alpha}}^i J_i|_N^P \kappa_{\underline{P}}^{\underline{\alpha}} \quad (4.3)$$

$$\kappa_{c\underline{\alpha}}^N = -ie^{i\phi} L^{NP} \epsilon_{\underline{\alpha}\underline{\beta}} \kappa_{P\underline{\beta}} \quad (4.4)$$

with $J_i|_N^P$ and L^{NP} some functions on the 5-sphere S_5 (they are properly defined in app.(B) where some of their useful properties are given).

We want now understand the relation between this vector field $\vec{\epsilon}$ and the associated variations $\delta_\kappa \Theta_N$, $\delta_\kappa \Theta_c^N$ and $\delta_\kappa x$. From eq. (4.3) we see clearly that the vector field $\vec{\epsilon}$ can be written as

$$\vec{\epsilon} = \epsilon_{\alpha N} \vec{D}^{\alpha N} \quad \psi_{\beta M} \left(\vec{D}^{\alpha N} \right) = \delta_\beta^\alpha \delta_M^N \quad (4.5)$$

and if we could expand $\vec{D}^{\alpha N}$ on the usual tangent basis $\partial^N = \frac{\partial}{\partial \Theta_N}$, $\partial_N = \frac{\partial}{\partial \Theta_c^N}$ and $\partial_x = \frac{\partial}{\partial x}$ then we could read the desired variations $\delta_\kappa \Theta_N$, $\delta_\kappa \Theta_c^N$ and $\delta_\kappa x$ since

$$\vec{\epsilon} = \delta_\kappa \Theta_N \vec{\partial}^N + \delta_\kappa \Theta_c^N \vec{\partial}_N + \delta_\kappa x \vec{\partial}_x \quad (4.6)$$

It is not very difficult to convince oneself that starting from a coset representative as in eq. (3.1) we can write

$$\delta_\kappa P \Theta_{\alpha N} = M_\alpha^\beta P \epsilon_{\beta N} + M_\alpha^\beta Q \epsilon_{\beta N} + N_\alpha^\beta Q \epsilon_{c\beta}^N + [O(\Xi^2)]_{\alpha N} \quad (4.7)$$

where M is a bosonic invertible matrix. The important point is that now we can set $Q\epsilon$ to zero and still have an invertible bosonic matrix connecting $P\epsilon_N$ and $\delta_\kappa P\Theta_N$: the fermionic correction does not matter since the bosonic part is invertible and hence it can be treated perturbatively.

In order to discuss the restrictions of the two projector types on the physical states which can be described we start noticing that using eq. (4.7) we can trade $P\epsilon_N$ for $\delta_\kappa P\Theta_N$, i.e. we can check when we can find the κ parameters (4.3) which allow to choose the gauge $P\epsilon_N = 0$ and then we can translate immediately the result to our actual κ variation $\delta_\kappa P\Theta_N$ because $P\epsilon_N$ and $\delta_\kappa P\Theta_N$ are connected by a non singular linear transformation. Obviously there will be some fermionic corrections but these can always be handled because the bosonic part is invertible. Hence the configurations where we cannot reach the gauge $P\epsilon_N = 0$ are the same where we cannot reach the true gauge $P\Theta_N = 0$. Because of this we can work with $\epsilon^{(+)} \equiv P\epsilon$ and then we can reread the results for $P\Theta$.

As a first step we use eq. (4.4) into eq. (4.3) to eliminate κ_{N1} then we write down the relation between $\epsilon^{(+)}$, its charge conjugate and the κ parameters for type I projectors

$$\epsilon_N^{(+)} = \mathcal{A}_0|_N^M \lambda_M + ie^{-i\phi} \mathcal{A}_1|_N^M L_{MP} \lambda_c^P + \mathcal{C}_0|_N^M \mu_M + ie^{-i\phi} \mathcal{C}_1|_N^M L_{MP} \mu_c^P \quad (4.8)$$

$$ie^{-i\phi} L_{NP} \epsilon_c^{P(+)} = \mathcal{A}_1|_N^M \lambda_M + ie^{-i\phi} \mathcal{A}_0|_N^M L_{MP} \lambda_c^P + \mathcal{C}_1|_N^M \mu_M + ie^{-i\phi} \mathcal{C}_0|_N^M L_{MP} \mu_c^P \quad (4.9)$$

and for type II projectors

$$\begin{aligned} \epsilon_N^{(+)} &= \mathcal{A}_0|_N^M \lambda_M + ie^{-i\phi} \mathcal{C}_1|_N^M L_{MP} \lambda_c^P + \\ &\quad + \mathcal{C}_0|_N^M \mu_M + ie^{-i\phi} \mathcal{A}_1|_N^M L_{MP} \mu_c^P \end{aligned} \quad (4.10)$$

$$\begin{aligned} ie^{-i\phi} L_{NP} \epsilon_c^{P(-)} &= \mathcal{C}_1|_N^M \lambda_M + ie^{-i\phi} \mathcal{A}_0|_N^M L_{MP} \lambda_c^P \\ &\quad + \mathcal{A}_1|_N^M \mu_M + ie^{-i\phi} \mathcal{C}_0|_N^M L_{MP} \mu_c^P \end{aligned} \quad (4.11)$$

where we have defined $\mu_M = P\kappa_{M0}$, $\lambda_M = Q\kappa_{M0}$ and \mathcal{A}, \mathcal{C} by means of the relations

$$\begin{aligned} QA = AP, PA = AQ &\implies \mathcal{A}_\alpha|_N^M = \Pi_\alpha^r \gamma_r \delta_N^M \\ QC = CQ, PC = CP &\implies \mathcal{C}_\alpha|_N^M = \Pi_\alpha^u \gamma_u \delta_N^M - i \Pi_\alpha^i J^i|_N^M \end{aligned}$$

where r runs on a proper subset R of the spacetime index set $\{0, 1, 2, 3, 4\}$ and u on its complement $\{0, 1, 2, 3, 4\} \setminus R$; both sets depend on the explicit form chosen for the matrix Γ entering the definition of the projector (2.1), for example

$$\Gamma = \gamma_0 \gamma_3 \implies R = \{0, 3\}$$

We can now make the following linearly independent combinations for type I projectors:

$$\begin{aligned} \epsilon_N^{(+)} \pm ie^{-i\phi} L_{NP} \epsilon_c^{P(+)} &= \mathcal{A}_\pm|_N^M [\lambda_M \pm ie^{-i\phi} L_{MP} \lambda_c^P] + \\ &\quad + \mathcal{C}_\pm|_N^M [\mu_M \pm ie^{-i\phi} L_{MP} \mu_c^P] \end{aligned} \quad (4.12)$$

and for type II projectors:

$$\begin{aligned} \epsilon_N^{(+)} \pm ie^{-i\phi} L_{NP} \epsilon_c^{P(-)} &= (\mathcal{A}_0 \pm \mathcal{C}_1)|_N^M [\lambda_M \pm ie^{-i\phi} L_{MP} \lambda_c^P] + \\ &\quad \pm (\mathcal{A}_1 \pm \mathcal{C}_0)|_N^M [\mu_M \pm ie^{-i\phi} L_{MP} \mu_c^P] \end{aligned} \quad (4.13)$$

Let us now discuss type I projectors only for brevity. We notice that the two equations in eq. (4.12) are linearly independent also with respect to the charge conjugation operation under which each equation transforms in itself, hence we can solve each of them independently of the other. As a first step we could try to solve the equations (4.12) for either $[\lambda_M \pm ie^{-i\phi} L_{MP} \lambda_c^P]$ or $[\mu_M \pm ie^{-i\phi} L_{MP} \mu_c^P]$. We find that in both cases the necessary condition for the non singularity of the matrices (\mathcal{A}_\pm) and (\mathcal{C}_\pm) is that

$$\Pi_\pm^r \Pi_{\pm r} = -\Pi_\pm^u \Pi_{\pm u} - \Pi_\pm^i \Pi_{\pm i} \neq 0 \quad (4.14)$$

where the equality is a consequence of the Virasoro constraints eq. (4.2). One could wonder whether this is a sufficient condition for the solvability of eq.s (4.12), the answer is yes since would the sum of the images of the two matrices (\mathcal{A}_\pm) and (\mathcal{C}_\pm) be the whole space spanned by Σ then it should be possible to find a number α such that $(\mathcal{A}_\pm + \alpha \mathcal{C}_\pm)$ is always invertible but it is easy to verify that the determinant of the previous combination is proportional to eq. (4.14) too.

We conclude therefore that states which do not satisfy eq. (4.14) cannot be described in this gauge.

A similar argument for type II projectors gives as in ([9])

$$\Pi_\tau^r \Pi_{\tau r} + \Pi_\sigma^u \Pi_{\sigma u} + \Pi_\sigma^i \Pi_{i\sigma} \neq 0 \quad (4.15)$$

5 The lightcone GS action for the type IIB superstring on $AdS_5 \times S_5$.

We want now to take the projector on the fermionic coordinate to be discarded to be

$$P = \frac{1}{\sqrt{2}} \gamma^0 \gamma^+ = \frac{1 + \gamma^0 \gamma^3}{2}$$

in such a way that we parametrize the κ gauge fixed (super)coset representative as

$$L = \exp \begin{pmatrix} & \theta_N \\ \bar{\theta}^N & \end{pmatrix} \exp \begin{pmatrix} i \frac{1+i\gamma^4}{2} \gamma_p x^p & \\ & 0_4 \end{pmatrix} \exp \begin{pmatrix} i \frac{\gamma^4}{2} \rho & \\ & 0_4 \end{pmatrix}$$

where we have defined

$$\theta_N = Q \Xi_N = \begin{pmatrix} \theta_{1N} \\ 0 \\ 0 \\ \theta_{2N} \end{pmatrix}$$

and ρ, x^p ($p = 0 \dots 3$) are the AdS_5 horospherical coordinates.

Starting from the usual formula for the coset geometry

$$L^{-1} dL = \begin{pmatrix} i \frac{1+i\gamma^4}{2} \gamma_p E^p & \Psi_N \\ \bar{\Psi}^N & 0_4 \end{pmatrix} + \text{spin-connection}$$

a tedious computation gives the $AdS_{5/4}$ potentials as

$$\begin{aligned} E^4 &= d\rho \\ E^v &= e^\rho dx^v \quad v = 1, 2, + \\ E^- &= e^\rho \left(dx^- + \frac{i}{2} \bar{\theta} \gamma^- d\theta \right) \\ \Psi_N &= e^{-\frac{i}{2} \gamma^4 \rho} \mathcal{M}^{-1} d\theta_N \quad \mathcal{M} = 1 + i \frac{1+i\gamma^4}{2} \gamma_p x^p \end{aligned}$$

in such a way that the fields entering the string action (4.1) can be written with the general formula of ([9]) in an explicit way as

$$\begin{aligned} E^- &= e^\rho \left(dx^- + \frac{i}{\sqrt{2}} \theta^{*Nn} d\theta_{Nn} \right) \\ A_+ &= -i \bar{\Psi}_{cN} \Psi_M \bar{\eta}_c^N \eta^M = 2e^\rho x^+ d\theta_{[N}^1 d\theta_{M]}^2 \bar{\eta}_c^N \eta^M \\ A_- &= -A_+^* \end{aligned}$$

where η^N ($N = 1 \dots 4$) are the 4 S_5 c-number killing spinors ([10])

$$\eta^N = \frac{1}{\sqrt{1-z^2}} (1 - z^i \tau_i) \epsilon^N$$

with ϵ^N are constant 5D-spinors which satisfy the normalization condition $\epsilon_N^\dagger \epsilon^M = \delta_N^M$ and τ^i are the corresponding 5D γ matrices and z^i are the S^5 projective coordinates .

In this way we can write the action in second order formalism as

$$\begin{aligned} S = \frac{T}{2} \int_{\Sigma_2} d^2\xi \sqrt{-g} g^{\alpha\beta} \frac{1}{2} \{ & \partial_\alpha x^\rho \partial_\beta x^\rho - e^{2\rho} [\partial_\alpha x^1 \partial_\beta x^1 + \partial_\alpha x^2 \partial_\beta x^2] \\ & + 2e^{2\rho} \partial_\alpha x^+ \left[\partial_\beta x^- + \frac{i}{\sqrt{2}} \theta^{*Nm} \overleftrightarrow{\partial}_\beta \theta_{Nm} \right] - 4\delta_{ij} \frac{\partial_\alpha z^i \partial_\beta z^j}{(1-z^2)^2} \} \\ & + \frac{e^{i\phi}}{2} e^\rho x^+ d\theta_{[N}^1 d\theta_{M]}^2 \bar{\eta}_c^N \eta^M - \frac{e^{-i\phi}}{2} e^\rho x^+ d\theta_1^{*[N} d\theta_2^{*M]} \bar{\eta}_N \eta_{cM} \end{aligned} \quad (5.1)$$

This action can be write in a more compact way if we introduce the 10D 32 components Majorana spinor defined by

$$\Theta = e^{-i\frac{\phi}{4}} \begin{pmatrix} e^{i\frac{\phi}{2}} \frac{1+i\gamma_4}{2} \theta_N \otimes \epsilon^N + e^{-i\frac{\phi}{2}} \frac{1-i\gamma_4}{2} \theta_c^N \otimes \epsilon_{cN} \\ -e^{-i\frac{\phi}{2}} \frac{1-i\gamma_4}{2} \theta_N \otimes \epsilon^N + e^{i\frac{\phi}{2}} \frac{1+i\gamma_4}{2} \theta_c^N \otimes \epsilon_{cN} \end{pmatrix} \quad (5.2)$$

and the coordinates

$$\begin{aligned} y^4 &= e^\rho \frac{1+z^2}{1-z^2} \\ y^i &= e^\rho \frac{2z^i}{1-z^2} \end{aligned} \quad (5.3)$$

so that we get

$$\begin{aligned} S = \frac{T}{2} \int_{\Sigma_2} d^2\xi \sqrt{-g} g^{\alpha\beta} \frac{1}{2} \{ & \frac{\partial_\alpha y^t \partial_\beta y_t}{y^2} \\ & - y^2 [\partial_\alpha x^1 \partial_\beta x^1 + \partial_\alpha x^2 \partial_\beta x^2] + 2y^2 \partial_\alpha x^+ [\partial_\beta x^- + \frac{i}{4} \bar{\Theta} \Gamma^- \partial_\beta \Theta] \\ & + x^+ y^t d\bar{\Theta} \Gamma_+ \Gamma_t d\Theta \end{aligned}$$

where $t = 4, 5, \dots 9$ is the transverse coordinates label as in ([3]).

This is not the end of the story since after the κ symmetry gauge fixing there should be 16 real fermionic d.o.f. while Θ has 32 real components: it is infact easy to verify that the spinor Θ (5.2) satisfies $\frac{1}{2} \Gamma^- \Gamma^+ \Theta = \Theta$. We can therefore change the Γ matrices representation to the usual lightcone basis in such a way that we can write the explicit form for the previous projector $\frac{1}{2} \Gamma^- \Gamma^+ = \begin{pmatrix} 1_{16} & \\ & 0_{16} \end{pmatrix}$ and then we can introduce a 8D spinor ζ so that

$$\Theta = \begin{pmatrix} \zeta \\ 0_{16} \end{pmatrix}$$

It is finally possible to take the flat limit rescaling $x^p \rightarrow e x^p, \rho \rightarrow e\rho, z^i \rightarrow e z^i, \Theta \rightarrow \sqrt{e} \Theta$ and letting $e \rightarrow 0$; we find that the previous action nicely reduces to the usual 10D flat lightcone action.

6 Conclusions.

In this letter we have classified the possible constant projector for the fixing of κ symmetry according to their behavior under Dirac conjugation and under charge conjugation and we have discussed the role of the different classes. It turns out that there is a preferred class, the S class (2.3), which always yields (super)vielbeins quadratic in the fermionic coordinates.

On the other hand the behaviour under charge conjugation (2.4,2.5) is connected to the constraint the physical states which can be described by the projector have to satisfy (4.14,4.15).

Using this knowledge we have then chosen the “simplest” gauge³, the one which corresponds to the lightcone in the flat limit, and we have constructed the GS action for the type IIB superstring on $AdS_5 \times S_5$ in this gauge. It turns nicely out that this action reduces to the usual GS action for the type IIB superstring in the flat limit. While the equations of motion cannot be solved exactly there is nevertheless hope that we can make use of this to start an effective large radius expansion as proposed in ([12]).

A $AdS_5 \otimes S_5$ conventions.

- Flat indices: $\widehat{a}, \widehat{b}, \dots \in \{0, \dots, 9\}$; $a, b, \dots \in \{0, 1, 2, 3, 4\}$, $i, j, \dots \in \{5, \dots, 9\}$; $p, q, \dots \in \{0, 1, 2, 3\}$, $t, u, v, \dots \in \{4, \dots, 9\}$; $\underline{\alpha}, \underline{\beta}, \dots \in \{0, 1\}$
- Flat metrics: $\eta_{\widehat{a}\widehat{b}} = \text{diag}(+1, -1, \dots, -1)$, $\eta_{ab} = \text{diag}(+1, -1, -1, -1, -1)$, $\eta_{ij} = \text{diag}(-1, -1, -1, -1, -1)$; $\eta_{\alpha\beta} = \text{diag}(+1, -1)$
- Epsilons: $\epsilon_{0\dots 9} = \epsilon_{0\dots 4} = \epsilon_{5\dots 9} = 1$; $\epsilon_{01} = 1$
- Charge conjugate spinors: $\Theta_c = C\overline{\Theta}^T$
- 1+9D gamma matrices

$$\begin{aligned} \{\Gamma_{\widehat{a}}, \Gamma_{\widehat{b}}\} &= 2\eta_{\widehat{a}\widehat{b}} \\ \Gamma_{11} &= \Gamma_0 \dots \Gamma_9 = \begin{pmatrix} 1_{16} & \\ & -1_{16} \end{pmatrix} \\ \Gamma_{\widehat{a}}^T &= -\widehat{C}^{-1} \Gamma_{\widehat{a}} \widehat{C} \quad \widehat{C}^T = -\widehat{C} \quad \widehat{C}^\dagger = \widehat{C}^{-1} \end{aligned}$$

- 1+4D (AdS_5) gamma matrices

$$\begin{aligned} \{\gamma_a, \gamma_b\} &= 2\eta_{ab} \\ \gamma_0 \gamma_1 \gamma_2 \gamma_3 \gamma_4 &= 1_4 \\ \gamma_a^T &= +C^{-1} \gamma_a C \quad C^T = -C \quad C^\dagger = C^{-1} \end{aligned}$$

Moreover we use the following (1+4)D γ explicit representation

³ As a matter of facts there is another gauge fixing with the same simplicity and it is given by $P = \frac{1+\gamma^0\gamma^4}{2}$

$$\gamma_p = \begin{pmatrix} \widetilde{\sigma}_p & \sigma_p \end{pmatrix} \gamma_4 = \begin{pmatrix} i \mathbf{1}_2 & \\ & -i \mathbf{1}_2 \end{pmatrix} C = \begin{pmatrix} i\sigma_2 & \\ & i\sigma_2 \end{pmatrix}$$

with $p, q, \dots = 0 \dots 3$, $\sigma_p = \{1_2, -\sigma_1, -\sigma_2, -\sigma_3\}$ and $\widetilde{\sigma}_p = \{1_2, \sigma_1, \sigma_2, \sigma_3\}$

- 0+5D (S_5) gamma matrices

$$\begin{aligned} \{\tau_i, \tau_j\} &= 2\eta_{ij} \\ \tau_5 \tau_6 \tau_7 \tau_8 \tau_9 &= i \mathbf{1}_2 \\ \tau_i^T &= +C^{-1} \tau_i C \quad C_5 = C \end{aligned}$$

- 1+9D gamma expressed through AdS_5 γ^a and S_5 τ^i gammas

$$\begin{aligned} \Gamma^{\widehat{a}} &= \{\gamma^a \otimes \mathbf{1}_4 \otimes \sigma_1, \mathbf{1}_4 \otimes \tau^i \otimes (-\sigma_2)\} \\ \widehat{C} &= C \otimes C_5 \otimes \sigma_2 \end{aligned}$$

Moreover we use the following (1+4)D γ explicit representation

$$\gamma_p = \begin{pmatrix} \widetilde{\sigma}_p & \sigma_p \end{pmatrix} \gamma_4 = \begin{pmatrix} i \mathbf{1}_2 & \\ & -i \mathbf{1}_2 \end{pmatrix} C = \begin{pmatrix} i\sigma_2 & \\ & i\sigma_2 \end{pmatrix}$$

with $p, q, \dots = 0 \dots 3$, $\sigma_p = \{1_2, -\sigma_1, -\sigma_2, -\sigma_3\}$ and $\widetilde{\sigma}_p = \{1_2, \sigma_1, \sigma_2, \sigma_3\}$

- 0+5D (S_5) gamma matrices

$$\begin{aligned} \{\tau_i, \tau_j\} &= 2\eta_{ij} \\ \tau_5 \tau_6 \tau_7 \tau_8 \tau_9 &= i \mathbf{1}_2 \\ \tau_i^T &= +C^{-1} \tau_i C \quad C_5 = C \end{aligned}$$

- 1+9D gamma expressed through AdS_5 γ^a and S_5 τ^i gammas

$$\begin{aligned} \Gamma^{\widehat{a}} &= \{\gamma^a \otimes \mathbf{1}_4 \otimes \sigma_1, \mathbf{1}_4 \otimes \tau^i \otimes (-\sigma_2)\} \\ \widehat{C} &= C \otimes C_5 \otimes \sigma_2 \end{aligned}$$

B Spinorial currents and useful Fierz identities.

We define the following useful quantities

$$J^i|_N^M = \overline{\eta}_N \tau^i \eta^M = \overline{\eta}_c^M \tau^i \eta_{cN} \quad (\text{B.1})$$

$$J^i|^{MN} = \overline{\eta}_c^N \tau^i \eta^M = -J^i|^{NM} \quad (\text{B.2})$$

$$J^i|_{MN} = \overline{\eta}_M \tau^i \eta_{cN} = -J^i|_{NM} \quad (\text{B.3})$$

$$L^{MN} = \overline{\eta}_c^M \eta^N = -L^{NM} \quad (\text{B.4})$$

$$L_{MN} = \overline{\eta}_M \eta_{cN} = -L_{NM} \quad (\text{B.5})$$

where η^N are the 4 S_5 c-number killing spinors ($N = 1 \dots 4$) defined by

$$D_{SO(5)}\eta^N \equiv \left(d - \frac{1}{4}\varpi^{ij}\tau_{ij} \right) \eta^N = -\frac{e}{2}\tau_i\eta^N E^i$$

In proving that the action (4.1) is invariant under the κ symmetry variations given by eq.s (4.3) it is important to use the following Fierz identities:

$$J^i|_N^M L^{NP} = L^{MP} J^i|_P^N = J^i|^{MN} \quad (\text{B.6})$$

$$L^{MN} L_{NP} = \delta_P^M \quad (\text{B.7})$$

$$J^i|_N^M J^j|_P^N + J^j|_N^M J^i|_P^N = 2\eta^{ij} \delta_P^M \quad (\text{B.8})$$

The derivation of these Fierz identities is based on the fact that η^N are a basis of the vector space on which the spinor irrep acts, i.e. $\bar{\eta}_N \tau \eta^N \propto \text{tr}(\tau)$.

References

- [1] R.R Metsaev and A.A. Tseytlin, Type IIB Superstring Action in $AdS_5 \times S_5$ Background, Nucl. Phys. B533 (1998) 109 (hep-th/9805028);
- [2] R. Kallosh, Superconformal Actions in Killing Gauge, hep-th/9807206
- [3] R. Kallosh and J. Rahmfeld, “The GS String Action on $AdS_5 \times S_5$ ”, Phys. Lett. B443 (1998) 143 (hep-th/9808038)
R. Kallosh and A.K. Tseytlin, “Simplifying Superstring Action on $AdS_5 \times S_5$ ”, JHEP 10(1998)016 (hep-th/9808088)
- [4] I. Pesando, The GS Type IIB Superstring Action on $AdS_3 \times S_3 \times T^4$, hep-th/9809145
- [5] J. Rahmfeld and A. Rajaraman, The GS String Action on $AdS_3 \times S^3$ with Ramond-Ramond Charge, hep-th/9809164
J. Park and S. Rey, Green-Schwarz Superstring on $AdS_3 \times S^3$, hep-th/9812062
- [6] P. Claus, “Super M-Branes Actions in $AdS_4 \times S^7$ and $AdS_7 \times S^4$ ”, hep-th/9809045
- [7] G. Dall’Agata, D. Fabbri, C. Fraser, P. Fre’, P. Termonia and M. Trigiante, The $Osp(8|4)$ Singleton Action from the Supermembrane, hep-th/9807115
- [8] P. Pasti, D. Sorokin and M. Tonin, “On Gauge-fixed Superbrane Actions in AdS Superbackgrounds”, hep-th/9809213
- [9] I. Pesando, “A κ Gauge Fixed Type IIB Superstring Action on $AdS_5 \times S_5$ ”, JHEP 11 (1998) 002 (hep-th/9808020)
- [10] I. Pesando, “All roads lead to Rome: Supersolvables and Supercosets”, hep-th/9808146

- [11] R.R Metsaev, “Light Cone Form of Field Dynamics in Anti-de Sitter Spacetime and AdS/CFT Correspondance”, hep-th/9906217
“Light Cone Gauge Formulation of IIB Supergravity in $AdS_5 \times S_5$ Background and AdS/CFT Correspondance”, hep-th/9908114
- [12] A. Rajaraman and M. Rozali, “On the Quantization of the GS String on $AdS_5 \times S_5$ ”, hep-th/9902046